

THIACALIX[4]ARENES DERIVATIVES: FROM SELF-ASSEMBLY TO FUNCTIONAL MATERIALS


I. S. Antipin,^{1,2} **S. E. Solovieva**,^{1,2} **V. A. Burilov**¹

¹Kazan Federal University, 420111, Russian Federation, Kazan, 18 Kremlevskaya St.

²A.E. Arbuzov Institute of Organic and Physical Chemistry, FRC Kazan Scientific Center, Russian Academy of Sciences, 420088, Russian Federation, Kazan, 8 Arbuzov St.

E-mail:iantipin54@yandex.ru

Abstract. Lower rim thiacalix[4]arenes derivatives in *cone* and *1,3-alternate* conformations have many advantages to create a wide range of precursors for the design of very sophisticated supramolecular architectures.



These compounds can be considered as technological platform for nanomaterials design by selfassembly method. Particular attention will be paid to the application of calixarene derivatives for the construction of various supramolecular and nanosystems, devices and “smart” materials: nanoparticles, Langmuir-Blodgett nanolayers, catalysts etc.

References

1. Thiacalix[4]arene-functionalized vesicles as phosphorescent indicators for pyridoxine detection in aqueous solution / V.A. Burilov, D.A. Mironova, R.R. Ibragimova, [et al.] // RSC Adv. – 2015.- Vol. 5. - P. 101177 - 101185.
 2. “Clickable” thiacalix[4]arene derivatives bearing polymerizable 1,3-butadiyne fragments: synthesis and incorporation into polydiacetylene vesicles / V. Burilov, A. Valiyakhmetova, D.Mironova [et al.] // RSC Adv. – 2016. – Vol. 6. - P. 44873-44877.
 3. Langmuir monolayers and thin films of amphiphilic thiacalix[4]arenes. properties and matrix for the immobilization of cytochrome c /S.E. Solovieva, R.A. Safiullin, E.N. Kochetkov [et al.] Langmuir. -2014. – Vol. 30. - P. 15153–15161.
 4. Novel amphiphilic conjugates of p-tert-butylthiacalix[4]arene with 10,12-pentacosadiynoic acid in 1,3-alternate stereoisomeric form. Synthesis and chromatic properties in the presence of metal ions. / V. Burilov, A. Valiyakhmetova, D. Mironova[et al.] // New J. Chem. - 2018. – Vol. 42. - P. 2942-2951.
- [95] Synthesis and binding with nucleoside phosphates of new polyammonium triazolyl amphiphiles on the base of p-tert-butylcalix[4]arene /V.A. Burilov, G.A. Fatikhova, M.N. Dokuchaeva, [et al.] // Beilstein J. Org. Chem. 2018. – Vol. 14. - P. 1980–1993.

This work was supported by the by Russian Science Foundation, project # 19-13-00095.